ELSEVIER

International Journal of Mass Spectrometry 231 (2004) 213

International Journal of Mass Spectrometry

www.elsevier.com/locate/ijms

Author index Volume 231

Abraham, G.T., 1
Abramov, S.V., 31
Afonso, C., 207
Al-Khalili, A., 37
Andrews, C.L., 169
Audier, H.E., 83, 197

Ballenthin, J.O., 17
Blais, J.-C., vi
Borshchevsky, A.Ya., 31
Boué, S.M., 147

Chatterjee, A., 1
Chilingarov, N.S., 31
Chou, C.-W., 147
Cole, R.B., 147, vi
Collado, V.M., 51
Cooks, R.G., 103
Cotter, R.J., ix

da Silveira, E.F., 51
De Pauw, E., 189
Debrauwer, L., 119
Dörner, R., 51
Du Fou de Kerdaniel, A., 131
Duursma, M., 37

Eberlin, M.N., 103

Farenzena, L., 51
Fenselau, C., 207
Flagan, R.C., 17
Frański, R., 47

Gabelica, V., 189
Gierczyk, B., 47
Giguère, M.-S., 59
Groeneveld, K.O., 51
Gronert, S., 179
Gross, M.L., 113
Guo, X., 37

Harsch, A., 169
Hathout, Y., 207
Hautreux, M., 131
Heeren, R.M.A., 37
Hori, K., 77
Huang, R., 179
Hue, N., 131

Jalowy, T., 51
Jouanin, I., 119

Kanno, H., 77
Karas, M., 189

Laprévote, O., 131
LeBlanc, B.W., 147
Li, K.H., 179
Li, Z., 69
Liu, S., 69

Malec, V., 131
March, R.E., 157
Mayer, P.M., 59
McDonnell, L.A., 37
Menon, A.G., 1
Meurer, E.C., 103
Miake-Lye, R.C., 17
Miao, X.-S., 157
Miller, T.M., 17
Morgan, L.R., 147
Mourgues, P., 83, 197

Nedev, H., 197

Page, A., 119
Paris, A., 119

Rathahao, E., 119
Rodgers, A.H., 147

Salcedo, D., 17
Schmidt-Böcking, H., 51
Schröder, D., 139
Schroeder, G., 47
Schwarz, H., 139
Seinfeld, J.H., 17
Sidorov, L.N., 31
Struck, R.F., 147

Thorn, W.F., 17
Timofeev, O., 113

van der Rest, G., 83, 197
Varutbangkul, V., 17
Viggiano, A.A., 17
Villalta, P.W., 17
Vouros, P., 169

Weber, Th., 51
Wormhoudt, J.C., 17
Worsnop, D.R., 17
Wu, L., 103

Yamada, Y., 77
Yang, P., 103
Young, B., 103

Zahir, A., 131
Zhu, M.M., 113

doi:10.1016/S1387-3806(04)00028-4

ELSEVIER

International Journal of Mass Spectrometry 231 (2004) 215–217

International Journal of Mass Spectrometry

www.elsevier.com/locate/ijms

Subject index Volume 231

Ab initio calculations
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83
Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197
Accurate mass
A fragmentation study of kaempferol using electrospray quadrupoletime-of-flight mass spectrometry at high mass resolution, 157
Acetone
Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69
N-Acetoxy-2-acetylaminofluorene
Analysis of the in vitro digestion of modified DNA to oligonucleotides by LC–MS and LC–MS/MS, 169
Aluminium fluoride
Mass spectrometric determination of partial pressures of ions in the saturated vapor over the $NaF–Na_3AlF_6$ system, 31
Averaging
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1

Binary cluster ion
Gas-phase structures of binary cluster ions of 1,4-dioxane and water, 77
BINOL
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139

CAD
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179
Carboxylate groups
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113
Catalyzed isomerizations
Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83
Catechol estrogens
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119
Central ion effects
Isomeric differentiation and quantification of α, β-amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103
CH bond activation
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139
Charge state
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131
Chemical ionization mass spectrometer
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17
CID
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
Collisionally induced dissociation
Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207
Collision-induced dissociation
Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69
Complexes
Isomeric differentiation and quantification of α, β-amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103
Cryolite
Mass spectrometric determination of partial pressures of ions in the saturated vapor over the $NaF–Na_3AlF_6$ system, 31

Density functional theory
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
Diastereoselectivity
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139
Dimer
Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207
1,4-Dioxane
Gas-phase structures of binary cluster ions of 1,4-dioxane and water, 77
Dissociation
Isomeric differentiation and quantification of α, β-amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103
Distonic ions
Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197
Distribution
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131
Disubstituted benzene
Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69

Electrospray
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131
Influence of the capillary temperature and the source pressure on the internal energy distribution of electrosprayed ions, 189

doi:10.1016/S1387-3806(04)00029-6

Electrospray ionization
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119
A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution, 157
Electrospray-ionization mass spectrometry
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
Enantioselectivity
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139
Energy resolved mass spectrometry
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119
Enzymatic digestion
Analysis of the in vitro digestion of modified DNA to oligonucleotides by LC–MS and LC–MS/MS, 169
Escape velocity
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1
ESI-MS
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113

Flavonoid
A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution, 157
FT-ICR mass spectrometry
Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83
Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197
FT-ICR-MS
Design and performance of a new FT-ICR cell operating at a temperature range of 77–438 K, 37

Gas phase derivatization
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179
Geometry factor
Design and performance of a new FT-ICR cell operating at a temperature range of 77–438 K, 37
Glutathione conjugates
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119

Heated capillary
Influence of the capillary temperature and the source pressure on the internal energy distribution of electrosprayed ions, 189
Hydrogen abstraction
Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197

Initial velocity
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51
Internal energy
Influence of the capillary temperature and the source pressure on the internal energy distribution of electrosprayed ions, 189
Ion dynamics
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1
Ion pressures
Mass spectrometric determination of partial pressures of ions in the saturated vapor over the $NaF–Na_3AlF_6$ system, 31
Ion trap mass spectrometry
Liquid chromatography coupled to negative electrospray/ion trap mass spectrometry for the identification of isomeric glutathione conjugates of catechol estrogens, 119
Ionized carbenes
Hydrogen radical abstraction by small ionized molecules, distonic ions and ionized carbenes in the gas phase, 197
Ionized dihydroxycarbenes
Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83
Ionized formic acid
Catalyzed isomerization and decarbonylation of ionized formic acid and dihydroxycarbene, 83
Ion–molecule equilibrium
Mass spectrometric determination of partial pressures of ions in the saturated vapor over the $NaF–Na_3AlF_6$ system, 31
Ion–molecule reaction
Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69
IPM
Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramide mustard, 147
Isomeric tripeptides
Isomeric differentiation and quantification of α, β-amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103
Isophosphoramide mustard
Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramide mustard, 147

Kaempferol
A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution, 157
Kinetic method
Isomeric differentiation and quantification of α, β-amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103

LC–ES–MS/MS
Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramide mustard, 147
Liquid chromatography–tandem mass spectrometry
Analysis of the in vitro digestion of modified DNA to oligonucleotides by LC–MS and LC–MS/MS, 169
Low temperature mass spectrometry
Design and performance of a new FT-ICR cell operating at a temperature range of 77–438 K, 37

Manganese complexes
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139
Mass spectrometric fragmentation pathway
Mass spectrometric fragmentation pathways of isotope labeled 2,5-disubstituted-1,3,4-oxadiazoles and thiadiazoles, 47
Mass spectrometry
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139

Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramide mustard, 147
Metallothionein
Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207
MM/MD
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59

Nanospray
Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207
Non-denaturing conditions
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131
Nonlinear Mathieu equation
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1

Oligonucleotide adducts
Analysis of the in vitro digestion of modified DNA to oligonucleotides by LC–MS and LC–MS/MS, 169
1,3,4-Oxadiazoles
Mass spectrometric fragmentation pathways of isotope labeled 2,5-disubstituted-1,3,4-oxadiazoles and thiadiazoles, 47

Paul trap
Escape velocity and resonant ion dynamics in Paul trap mass spectrometers, 1
Peptide analysis
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179
Peptides
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113
Perchloric acid
Gas chromatography–electron ionization mass spectrometry and liquid chromatography–electrospray tandem mass spectrometry for determination of impurities in the anti-cancer drug isophosphoramide mustard, 147
Phosphopeptides
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179
Poly(vinyl acetate)
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59
Position sensitive detector
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51
Pressure effect
Influence of the capillary temperature and the source pressure on the internal energy distribution of electrosprayed ions, 189
Proteins
Under non-denaturing solvent conditions, the mean charge state of a multiply charged protein ion formed by electrospray is linearly correlated with the macromolecular surface, 131
Proteomics
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113

Quantification
Isomeric differentiation and quantification of α, β-amino acid-containing tripeptides by the kinetic method: alkali metal-bound dimeric cluster ions, 103

Relative humidity
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17

Secondary ion desorption
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51
Skeletal rearrangement
Mass spectrometric fragmentation pathways of isotope labeled 2,5-disubstituted-1,3,4-oxadiazoles and thiadiazoles, 47
Sodium ion interactions
Information for proteomics: ESI-MS titration by sodium ions gives the number of carboxylate groups in peptides, 113
Substituent effect
Substituent effect on ion–molecule reaction of disubstituted benzenes with acetone plasma, 69
Sulfur dioxide
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17
Sulfuric acid
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17

Tandem mass spectrometry
A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution, 157
Temperature-controlled ICR cell
Design and performance of a new FT-ICR cell operating at a temperature range of 77–438 K, 37
1,3,4-Thiadiazoles
Mass spectrometric fragmentation pathways of isotope labeled 2,5-disubstituted-1,3,4-oxadiazoles and thiadiazoles, 47
Time-of-flight
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51
Time-of-flight mass spectrometry
Gas-phase structures of binary cluster ions of 1,4-dioxane and water, 77
Trimethyl borate
Gas phase derivatization in peptide analysis I: the utility of trimethyl borate in identifying phosphorylation sites, 179

Unimolecular decomposition
Climbing the internal energy ladder: the unimolecular decomposition of ionized poly(vinyl acetate), 59

Vanadium complexes
Diastereo- and enantioselective bond activation of alkanols by gaseous metal ions, 139

Water
Gas-phase structures of binary cluster ions of 1,4-dioxane and water, 77
Water clusters
Effect of relative humidity on the detection of sulfur dioxide and sulfuric acid using a chemical ionization mass spectrometer, 17

XY-TOF
Initial velocity of secondary ions from *XY*-TOF technique, simultaneous calibration by residual gas ionization, 51

Zinc chelates
Evidence for zinc ion sharing in metallothionein dimers provided by collision-induced dissociation, 207